



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Hypoxyprobe™ reveals dynamic spatial and temporal changes in hypoxia in a mouse model of endometrial breakdown and repair

Citation for published version:

Cousins, F, Murray, A, Scanlon, JP & Saunders, P 2016, 'Hypoxyprobe™ reveals dynamic spatial and temporal changes in hypoxia in a mouse model of endometrial breakdown and repair', *BMC Research Notes*. <https://doi.org/10.1186/s13104-016-1842-8>

Digital Object Identifier (DOI):

[10.1186/s13104-016-1842-8](https://doi.org/10.1186/s13104-016-1842-8)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

BMC Research Notes

Publisher Rights Statement:

© 2016 Cousins et al. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



CASE REPORT

Open Access



HypoxyprobeTM reveals dynamic spatial and temporal changes in hypoxia in a mouse model of endometrial breakdown and repair

Fiona L. Cousins¹, Alison A. Murray², Jessica P. Scanlon¹ and Philippa T. K. Saunders^{1*} 

Abstract

Background: Menstruation is the culmination of a cascade of events, triggered by the withdrawal of progesterone at the end of the menstrual cycle. Initiation of tissue destruction and endometrial shedding causes spiral arteriole constriction in the functional layer of the endometrium. Upregulation of genes involved in angiogenesis and immune cell recruitment, two processes that are essential to successful repair and remodelling of the endometrium, both thought to be induced by reduced oxygen has been reported. Evidence for stabilisation/increased expression of the transcriptional regulator hypoxia inducible factor in the human endometrium at menses has been published. The current literature debates whether hypoxia plays an essential role during menstrual repair, therefore this study aims to delineate a role for hypoxia using a sensitive detection method (the HypoxyprobeTM) in combination with an established mouse model of endometrial breakdown and repair.

Results: Using our mouse model of menses, during which documented breakdown and synchronous repair occurs in a 24 h timeframe, in combination with the HypoxyprobeTM detection system, oxygen tensions within the uterus were measured. Immunostaining revealed striking spatial and temporal fluctuations in hypoxia during breakdown and showed that the epithelium is also exposed to hypoxic conditions during the repair phase. Furthermore, time-dependent changes in tissue hypoxia correlated with the regulation of mRNAs encoding for the angiogenic genes vascular endothelial growth factor and stromal derived factor (*Cxcl12*).

Conclusions: Our findings are consistent with a role for focal hypoxia during endometrial breakdown in regulating gene expression during menses. These data have implications for treatment of endometrial pathologies such as heavy menstrual bleeding.

Keywords: Endometrium, Hypoxia, Menstruation, Repair, Angiogenesis

Background

In women demise of the corpus luteum in the absence of pregnancy results in a rapid decline in peripheral progesterone that precipitates a cascade of molecular changes which induce myometrial contractions and vasoconstriction of the spiral arterioles of the endometrium. The net result of these changes is the generation of a hypoxic microenvironment in the upper, decidualised, functional

layer of the endometrium which is typical of that associated with tissue necrosis and ischaemic injury [1]. Under normoxia, cellular oxygen sensors hydroxylate specific proline residues in the HIF-1 α (hypoxia inducible factor-1 α) protein targeting it for rapid proteasome dependent degradation [2, 3]. Under hypoxic conditions, degradation of HIF-1 α does not occur and the protein translocates to the nucleus where it forms heterodimeric binding complexes with HIF- β [4]. The HIF heterodimer binds to hypoxic response elements, acting as a transcription factor regulating expression of hypoxic response genes a number of which are expressed in the endometrium during menses [5–7].

*Correspondence: p.saunders@ed.ac.uk

¹ Medical Research Council Centre for Inflammation Research, The University of Edinburgh, 47 Little France Crescent, Edinburgh EH16 4TJ, UK

Full list of author information is available at the end of the article

HIF1 α protein and its mRNA have been identified in the human endometrium, however the number of samples recovered during the menstrual phase was low with only $n = 2$ [5]. Previous studies using HIF expression as a surrogate for detection of hypoxia in the endometrium have yielded inconsistent results [5, 8]. As HIF is notoriously unstable it is possible that expression is lost during tissue collection and fixation or that the duration of hypoxia was too brief to stabilise enough HIF for immunohistochemical detection.

There is some disagreement in the literature as to whether hypoxia plays an essential role in regulation of tissue breakdown and initiation of repair during menses. Notably it is reported that the pattern of immunorepression of HIF1 α , HIF1 β and HIF2 α across the menstrual cycle is inconsistent with a role for hypoxia at menses [8]. Conversely in the same study, in vitro culture of endometrial stromal cells under hypoxic conditions resulted in an increase in intracellular HIF1 α and VEGF [8] and together with data from Maybin et al. [9] would appear to support a role for hypoxia at menses.

These apparently contradictory results have prompted use of alternatives to immunohistochemistry including electron paramagnetic resonance (EPR) oximetry in combination with implanted particulate materials for in vivo measurement of oxygen tension. However, the use of these particulates does not generate images of the sample sites and successful EPR is highly dependent upon the microenvironment surrounding the particulate. A xenograft model of menstruation has utilised EPR in combination with lithium phtalocyanine (LiPc) crystals, which are sensitive to oxygen levels below 10 mmHg [10], this study reported no consistent evidence of hypoxia during menses. However, in this study the tissue was restricted to samples from the functional layer and therefore lacked the spiral arterioles of the basal layer that are thought to vasoconstrict at menstruation. An additional complication was that the model did not consider the role that the microenvironment of the flank engraftment site may play in xenograft establishment and HIF degradation during steroid induced 'menstruation' in the human tissue fragment.

The development of the Hypoxyprobe™ system facilitated direct visualisation of hypoxia in situ at oxygen concentrations of <1 %, irrespective of how long a cell has been under hypoxic conditions, but more importantly detection of the probe is resistant to fixation. The Hypoxyprobe™ system uses pimonidazole hydrochloride as its hypoxia marker, which forms protein adducts in hypoxic cells ($pO_2 < 10$ mmHg) when administered in vivo. These protein adducts can be visualised by a specific monoclonal antibody. Detection of pimonidazole has been shown to be a reliable and direct measure of tissue hypoxia.

We hypothesised that hypoxia does occur during endometrial breakdown and that it is therefore likely to be a major player in promoting changes in gene expression that support rapid tissue repair at menses. Using the Hypoxyprobe™ in conjunction with a validated mouse model of rapid endometrial breakdown and repair revealed a spatial and temporal gradient of hypoxia in the endometrium which was associated with temporal changes in expression of angiogenic factors that are integral to restoration of a normal endometrial architecture.

Methods

C57BL6 mice of 8–12 weeks of age were used in a previously validated protocol [11] in combination with the Hypoxyprobe™ labelling system. Experiments were performed in accordance with UK legal requirements and under licensed approval from the UK Home Office (London) following Ethical Review. Mice were given access to food and water ad libitum and housed in accordance with Home Office guidelines. Briefly, mice underwent the menses protocol as detailed previously. Mice were culled at specific time-points; 0 h (time of progesterone withdrawal, achieved by the removal of a progesterone-secreting pellet, PWD), 4 h (after PWD), 8 h (after PWD) and 24 h (after PWD). Ninety minutes prior to culling each mouse received a single intra-peritoneal injection of the pimonidazole (15 mg/ml in PBS). Uteri were dissected, fixed in 4 % neutral buffered formalin and processed into paraffin wax. Hypoxia was investigated in transverse uterine sections using standard protocols facilitating detection of pimonidazole with the Hypoxyprobe™ antibody (diluted 1/100 in normal animal serum). Hypoxia, defined as <1 % oxygen was visualised as a brown stain following immunohistochemistry by DAB chromogen detection. QRTPCR analysis was performed as described previously [11]; *Vegfa* Fwd aaaaacgaagcgcaagaaa Rev ttctccgctctgaacaagg, *Cxcl12* Fwd ctgtgcccttcagattgtt Rev taatttcgggtcaatgcaca, *Flt1* Fwd ggcccgggatattataag aac Rev ccatccatttaggggaagtc, *Kdr* Fwd cagtgggtactggcagctagaag Rev acaagcatagcgggctgtt. Each sample was expressed relative to the endogenous control 18 s and then compared to the mean 0 h time-point sample set. Statistical analysis was performed by One Way ANOVA with Bonferroni post hoc testing where statistical significance was accepted where $p < 0.05$. Histograms for each gene are depicted as the mean \pm the standard error of the mean.

Results

Immunostaining for the Hypoxyprobe™ identified spatial and temporal hypoxia in the endometrium. At time of PWD (0 h, Fig. 1A) the endometrium was largely unstained for pimonidazole, an indication of normoxia.




Fig. 1 The endometrium is exposed to hypoxic conditions in a temporal and spatial manner during endometrial breakdown and repair. **A** Time of progesterone withdrawal (0 h), hypoxia is detected in the decidualised cell mass (inset **a**), **B** 4 h after progesterone withdrawal (PWD), the decidualised cell mass is positively stained for pimonidazole. The luminal epithelium is unstained (inset **b**, arrows), **C** The intensity of staining is stronger at 8 h after PWD. A distinct gradient is detected at the border of basal layer and the decidualised functional layer (inset **c**), **D** At 24 h after PWD the re-epithelialisation luminal epithelium and the adjacent stroma is immunopositive (inset **d**). BL basal layer, DC decidualised cells, SC shed cells, asterisk evidence of hemosiderin, NPC negative primary control

An area in the decida was observed to be positively stained for pimonidazole protein adducts (Fig. 1A, inset a). Hemosiderin, most commonly found in macrophages, was observed in the basal layer (Fig. 1A, asterisk). This observation is consistent with findings from Kaitu'u-Lino et al. who observed macrophages residing in the basal layer at the time of PWD [12]. Within 4 h of PWD (Fig. 1B) the decidualised cell mass was intensely stained for hypoxia, whilst the basal layer remained unstained. The luminal epithelium was negative for hypoxia (indicated by the arrows in Fig. 1B, inset b), whilst the stromal cells adjacent to the luminal epithelium were positively stained. The intensity of staining of pimonidazole adducts was stronger in the shedding decidual cell mass in tissues collected 8 h after the withdrawal of progesterone (Fig. 1C). The luminal epithelium remained unstained. The basal layer also remained normoxic, with the existence of a defined gradient in the intensity of staining in the decidualised cells and the underlying basal layer (Fig. 1C, inset c) consistent with spatial

differences in hypoxia within the tissue. Within 24 h the uterine horn had regressed in size and the shed cells in the lumen were largely unstained for hypoxia (Fig. 1D). However, the luminal epithelium and the adjacent stroma were intensely stained (Fig. 1D, inset d) indicating that the endometrium is still under hypoxic conditions during endometrial repair.

It has been reported that the genes implicated in regulation of angiogenesis including *Vegfa* (vascular endothelial growth factor a) and *Cxcl12* (also known as stromal derived factor) are regulated by hypoxia in human endometrial stromal cells in vitro [9, 13]. Treatment of mice with a potent Vegf blocker (Vegf-trap) blocks neovascularisation and inhibits re-epithelialisation in endometrium of both macaque and mouse [14]. In addition, VEGF receptor 1 (VEGFR1 also known as *Flt1*) and VEGFR2 (*Kdr*) have been reported to be differentially regulated by hypoxia in a number of organs and cell types [15–17], therefore the abundance of the mRNAs encoded by these genes was investigated by




Fig. 2 Onset of menstruation and hypoxia results in differential regulation of angiogenic genes *Vegfa*, *Cxcl12*, *Flt1* (VEGFR1) and *Kdr* (VEGFR2) All samples were normalised against the endogenous control 18 s then compared relative to the mean of 0 h time-point sample set. Statistical analysis was performed by one way ANOVA using Bonferroni post hoc testing, where statistical significance was accepted at $p < 0.05$. Histograms are depicted as the mean \pm the standard error of the mean

qRT-PCR at the same time points (Fig. 2). Concentrations of mRNAs for *Vegfa* were significantly increased at 8 h ($p < 0.001$) after progesterone withdrawal, consistent with intense immunostaining for hypoxia at this time-point. *Vegfa* mRNA concentrations were significantly decreased at 24 h ($p < 0.01$), when compared to the 8 h time-point. In contrast there was no change in mRNA concentrations for *Cxcl12* between 0 and 8 h, consistent with the tissue being under hypoxic conditions. As the tissue breaks down and the hypoxic decidualised cell mass is shed, a significant increase in *Cxcl12* was observed ($p < 0.001$) consistent with findings from Tsuzuki et al. [13] who observed a decrease in CXCL12 mRNA expression in human stromal cells

under hypoxia in comparison to cells incubated under normoxic conditions. In contrast to findings in the brain, kidney and liver where VEGFR1 was induced by hypoxia, *Flt1* mRNA concentrations were significantly down-regulated at 4 h ($p < 0.001$, Fig. 2) coincident with intense staining for hypoxia, *Flt1* mRNAs were still decreased, in comparison with the 0 h time-point, at the 24 h time-point suggesting that hypoxia does regulate VEGFR1 but in a tissue dependent manner. Consistent with findings by Coudyzer et al. [10] *Kdr* mRNA concentrations did not change after PWD and the onset of hypoxia (between the 0 and 8 h time-point) supporting data from other body organs which reports that VEGFR2 is not regulated by hypoxia [15].

Conclusions

Using the Hypoxyprobe™ in combination with a mouse model in which shedding and repair of endometrium within 24 h after progesterone withdrawal can be studied, intense immunostaining for hypoxia during breakdown and shedding of the decidual mass has been shown. Examination of tissue sections revealed that a gradient of hypoxia develops during repair of the tissue consistent with regulation of genes implicated in regulation of neo-angiogenesis and epithelial repair.

Abbreviations

ANOVA: analysis of variance; EPR: electron paramagnetic resonance; HIF/HIF1 α / β : hypoxia inducible factor/hypoxia inducible factor 1 α /beta; LiPc: lithium phthalocyanine crystals; pO₂: partial pressure of oxygen; PWD: progesterone withdrawal; VEGF(R): vascular endothelial growth factor (receptor).

Authors' contributions

FLC was involved in study design, data collection, data analysis and writing of manuscript. AAM was involved in data collection. JCS was involved in data analysis. PTKS was involved in study design, data analysis and writing of the manuscript. All animal work was performed under the project license held by PTKS. All authors read and approved the final manuscript.

Author details

¹ Medical Research Council Centre for Inflammation Research, The University of Edinburgh, 47 Little France Crescent, Edinburgh EH16 4TJ, UK. ² Medical Research Council Centre for Reproductive Health, The University of Edinburgh, 47 Little France Crescent, Edinburgh EH16 4TJ, UK.

Acknowledgements

The authors would like to acknowledge Phoebe Kirkwood for assisting with figure preparation and Dr Douglas Gibson for critical reading of the manuscript. The studies were funded by a MRC Grant G1100356/1 to PTKS and a MRC studentship for FLC.

Competing interests

The authors declare that they have no competing interests.

Received: 30 October 2015 Accepted: 7 January 2016

Published online: 19 January 2016

References

- Markee JE. Menstruation in intraocular endometrial transplants in the rhesus monkey. *Contrib Embryol.* 1940;28:219–318.
- Huang LE, Arany Z, Livingston DM, Bunn HF. Activation of hypoxia-inducible transcription factor depends primarily upon redox-sensitive stabilization of its subunit. *J Biol Chem.* 1996;271:32253–9. doi:10.1074/jbc.271.50.32253.
- Huang LE, Gu J, Schau M, Bunn HF. Regulation of hypoxia-inducible factor 1 α is mediated by an O₂-dependent degradation domain via the ubiquitin-proteasome pathway. *Proc Natl Acad Sci U S A.* 1998;95:7987–92.
- Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci USA.* 1995;92:5510–4.
- Critchley HOD, Osei J, Henderson TA, et al. Hypoxia-inducible factor-1 α expression in human endometrium and its regulation by prostaglandin E-series prostanoid receptor 2 (EP2). *Endocrinology.* 2006;147:744–53. doi:10.1210/en.2005-1153.
- Maybin JA, Hirani N, Jabbour HN, Critchley HOD. Novel roles for hypoxia and prostaglandin E2 in the regulation of IL-8 during endometrial repair. *Am J Pathol.* 2011;178:1245–56. doi:10.1016/j.ajpath.2010.11.070.
- Maybin JA, Barcroft J, Thiruchelvam U, et al. The presence and regulation of connective tissue growth factor in the human endometrium. *Hum Reprod.* 2012;27:1112–21. doi:10.1093/humrep/der476.
- Zhang J, Salamsen LA. Expression of hypoxia-inducible factors in human endometrium and suppression of matrix metalloproteinases under hypoxic conditions do not support a major role for hypoxia in regulating tissue breakdown at menstruation. *Hum Reprod.* 2002;17:265–74.
- Maybin JA, Hirani N, Brown P, et al. The regulation of vascular endothelial growth factor by hypoxia and prostaglandin F₂ α during human endometrial repair. *J Clin Endocrinol Metab.* 2011;96:2475–83. doi:10.1210/jc.2010-2971.
- Coudyzer P, Lemoine P, Jordan BF, et al. Hypoxia is not required for human endometrial breakdown or repair in a xenograft model of menstruation. *FASEB J.* 2013. doi:10.1096/fj.13-232074.
- Cousins FL, Murray A, Esnal A, et al. Evidence from a mouse model that epithelial cell migration and mesenchymal-epithelial transition contribute to rapid restoration of uterine tissue integrity during menstruation. *PLoS One.* 2014;9:e86378. doi:10.1371/journal.pone.0086378.
- Kaitu'u-Lino TJ, Morison NB, Salamsen LA. Neutrophil depletion retards endometrial repair in a mouse model. *Cell Tissue Res.* 2007;328:197–206. doi:10.1007/s00441-006-0358-2.
- Tsuzuki T, Okada H, Cho H, et al. Hypoxic stress simultaneously stimulates vascular endothelial growth factor via hypoxia-inducible factor-1 α and inhibits stromal cell-derived factor-1 in human endometrial stromal cells. *Hum Reprod.* 2012;27:523–30. doi:10.1093/humrep/der405.
- Fan X, Krieg S, Kuo CJ, et al. VEGF blockade inhibits angiogenesis and reepithelialization of endometrium. *FASEB J.* 2008;22:3571–80. doi:10.1096/fj.08-111401.
- Marti HH, Risau W. Systemic hypoxia changes the organ-specific distribution of vascular endothelial growth factor and its receptors. *Proc Natl Acad Sci.* 1998;95:15809–14. doi:10.1073/pnas.95.26.15809.
- Okuyama H, Krishnamachary B, Zhou YF, et al. Expression of vascular endothelial growth factor receptor 1 in bone marrow-derived mesenchymal cells is dependent on hypoxia-inducible factor 1. *J Biol Chem.* 2006;281:15554–63. doi:10.1074/jbc.M602003200.
- Ulyatt C, Walker J, Ponnambalam S. Hypoxia differentially regulates VEGFR1 and VEGFR2 levels and alters intracellular signaling and cell migration in endothelial cells. *Biochem Biophys Res Commun.* 2011;404:774–9. doi:10.1016/j.bbrc.2010.12.057.

Submit your next manuscript to BioMed Central and we will help you at every step:

- We accept pre-submission inquiries
- Our selector tool helps you to find the most relevant journal
- We provide round the clock customer support
- Convenient online submission
- Thorough peer review
- Inclusion in PubMed and all major indexing services
- Maximum visibility for your research

Submit your manuscript at
www.biomedcentral.com/submit

